


ADVATE
[Antihemophilic Factor
(Recombinant)]


ADYNOVATE
[Antihemophilic Factor
(Recombinant), PEGylated]



KEEP PAINTING

YOUR STORY

Join us for an interactive program to help you define your goals and create a picture of where you want to go next. Discover more about hemophilia A and becoming the artist of your own journey.

This program includes information about ADVATE® and ADYNOVATE® treatment options presented by a Takeda Community Education Specialist (CES).

DATE: Thursday, February 27, 2025

TIME: 6:00 PM Central Standard Time

LOCATION: Tiphonie's, 210 East 2nd Street, Davenport, IA

RSVP: Brooke Loving: bdh_director@bdheartland.org, 319-721-1664

SPEAKER: Kevin Ellebrecht

Attendees and Guests

This program is for U.S. patients and caregivers only and is not intended for healthcare professionals. All patient attendees must be of an appropriate age for the disease state and/or products being discussed. Patient attendees may bring up to two caregivers/family members. Attendees under the age of 18 must register with and be accompanied by an adult. Attendee expenses are not reimbursable. This program is for educational purposes only and is not a substitute for the advice provided by your healthcare provider. Consult your healthcare professional for your medical concerns.

IMPORTANT INFORMATION

What is ADYNOVATE and ADVATE?

- ADYNOVATE and ADVATE are each prescription, injectable medicines that are used to replace clotting factor, to help treat and control bleeding in children and adults with hemophilia A (congenital Factor VIII deficiency, also called “classic” hemophilia).
 - Your healthcare provider (HCP) may give you ADYNOVATE or ADVATE when you have surgery.
 - ADYNOVATE and ADVATE can each reduce the number of bleeding episodes when used regularly (prophylaxis).
- ADYNOVATE and ADVATE are not used to treat von Willebrand disease.

DETAILED IMPORTANT RISK INFORMATION

Who should not use ADYNOVATE and ADVATE?

Do not use ADYNOVATE or ADVATE if you:

- Are allergic to mouse or hamster proteins.
- Are allergic to any ingredients in ADYNOVATE or ADVATE.

Tell your HCP if you are pregnant or breastfeeding because ADYNOVATE or ADVATE may not be right for you.

Please see additional safety information on back, and enclosed Prescribing Information for ADVATE and ADYNOVATE located in pocket and discuss with your HCP.



DETAILED IMPORTANT RISK INFORMATION (continued)

What should I tell my HCP before using ADYNOVATE and ADVATE?

Tell your HCP if you:

- Have or have had any medical problems.
- Take any medicines, including prescription and non-prescription medicines, such as over-the-counter medicines, supplements or herbal remedies.
- Have any allergies, including allergies to mice or hamster proteins.
- Are breastfeeding. It is not known if ADYNOVATE or ADVATE pass into your milk or if they can harm your baby.
- Are or become pregnant. It is not known if ADYNOVATE or ADVATE may harm your unborn baby.
- Have been told that you have inhibitors to factor VIII (because ADYNOVATE or ADVATE may not work for you).

What important information do I need to know about ADYNOVATE and ADVATE?

- You can have an allergic reaction to ADYNOVATE or ADVATE. Call your healthcare provider right away and stop treatment if you get a rash or hives, itching, tightness of the throat, chest pain or tightness, difficulty breathing, lightheadedness, dizziness, nausea or fainting.
- Do not attempt to infuse yourself with ADYNOVATE or ADVATE unless you have been taught by your HCP or hemophilia center.

What else should I know about ADYNOVATE, ADVATE, and Hemophilia A?

- Your body may form inhibitors to factor VIII. An inhibitor is part of the body's normal defense system. If you form inhibitors, it may stop ADYNOVATE or ADVATE from working properly. Talk with your HCP to make sure you are carefully monitored with blood tests for the development of inhibitors to factor VIII.

What are possible side effects of ADYNOVATE and ADVATE?

- **ADYNOVATE:** The common side effects of ADYNOVATE are headache, diarrhea, rash, nausea, dizziness, and hives. These are not all the possible side effects with ADYNOVATE. Tell your HCP about any side effects that bother you or do not go away.
- **ADVATE:** Side effects that have been reported with ADVATE include: cough, headache, joint swelling/aching, sore throat, fever, itching, unusual taste, dizziness, hematoma, abdominal pain, hot flashes, swelling of legs, diarrhea, chills, runny nose/congestion, nausea/vomiting, sweating, and rash. Tell your HCP about any side effects that bother you or do not go away or if your bleeding does not stop after taking ADVATE.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

For additional safety information, please see the enclosed Prescribing Information for ADVATE, and also for ADYNOVATE located in the pocket and discuss with your HCP.